


IRON REDOX TITRATIONS



Name Form

- 24.3 cm³ of 0.02 mol dm⁻³ KMnO₄ reacted with 20.0 cm³ of an iron (II) solution. Calculate the molarity of the iron (II) ion.
- Calculate the percentage of iron in a sample of steel wire if 1.51 g of the wire was dissolved in excess of dilute sulphuric acid and the solution made up to 250 cm³ in a standard flask. 25.0 cm³ of this solution was pipetted into a conical flask and needed 25.45 cm³ of 0.02 mol dm⁻³ KMnO₄ for complete oxidation.
- 3.00 g of a lawn sand containing an iron (II) salt was shaken with dilute H₂SO₄. The resulting solution required 25.00 cm³ of 0.0200 mol dm⁻³ potassium manganate (VII) to oxidise the Fe²⁺ ions in the solution to Fe³⁺ ions. Use this to calculate the percentage by mass of Fe²⁺ ions in this sample of lawn sand.
- Calculate x in the formula FeSO₄.xH₂O from the following data: 12.18 g of iron (II) sulphate crystals were made up to 500 cm³ acidified with sulphuric acid. 25.0 cm³ of this solution required 43.85 cm³ of 0.01 mol dm⁻³ KMnO₄ for complete oxidation.
- A tablet weighing 0.940 g was dissolved in dilute sulphuric acid made up to 250 cm³ with water. 25.0 cm³ of this solution was titrated with 0.00160 M K₂Cr₂O₇ requiring 32.5 cm³ of the K₂Cr₂O₇. Calculate the percentage by mass of Fe²⁺ in the tablet.
- Calculate the volume of 0.0200 mol dm⁻³ potassium manganate (VII) which just reacts with 0.142 g of iron (II) sulphate, in acid solution.
- Ammonium iron (II) sulphate crystals have the following formula: (NH₄)₂SO₄.FeSO₄.nH₂O. In an experiment to find n, 8.492 g of the salt were dissolved and made up to 250 cm³ solution with distilled water and dilute sulphuric acid. A 25.0 cm³ portion of the solution was titrated against 0.0150 mol dm⁻³ KMnO₄, 22.5 cm³ being required. Calculate n.
- A piece of iron wire weighs 2.225 g. It is dissolved in acid, which oxidises it to Fe²⁺ and made up to 250 cm³. A 25 cm³ sample required 31.0 cm³ of a 0.0185 mol dm⁻³ solution of potassium dichromate. Calculate the percentage of iron in the wire.
- A 25.0 cm³ aliquot of a solution containing Fe²⁺ and Fe³⁺ ions was acidified and titrated against 0.0200 M potassium manganate (VII) solution, requiring 15.0 cm³. Zn reduces Fe³⁺ to Fe²⁺ and a second aliquot was reduced by zinc and after filtering off the excess zinc, was titrated with the same potassium manganate solution, requiring 19.0 cm³. Calculate the concentrations of the Fe²⁺ and Fe³⁺ in the solution.
- 13.2 g of iron (III) alum were dissolved in water and reduced to an iron(II) ion solution by zinc and dilute sulphuric acid. The mixture was filtered and the filtrate and washings made up to 500 cm³ in a standard volumetric flask. 20.0 cm³ of this solution required 26.5 cm³ of 0.01 mol dm⁻³ KMnO₄ for oxidation. Calculate the percentage by mass of iron in iron alum.
- A sample of solid ethanedioic acid (H₂C₂O₄.2H₂O) has been contaminated with potassium ethanedioate (K₂C₂O₄.xH₂O). A 1.780 g sample of this mixture was made up to a 250 cm³ solution with distilled water. A 25 cm³ sample was titrated against 0.100 mol dm⁻³ sodium hydroxide, requiring 17.35 cm³. Another 25 cm³ sample was acidified with sulphuric acid and titrated against 0.0200 mol dm⁻³ KMnO₄ solution, requiring 24.85 cm³. Calculate x.

$$\text{C}_2\text{O}_4^{2-} \rightarrow 2 \text{CO}_2 + 2 \text{e}^-$$
- A piece of rusted iron was analysed to find out how much of the iron had been oxidised to rust [hydrated iron(III) oxide]. A small sample of the iron was dissolved in excess dilute sulphuric acid to give 250 cm³ of solution. The solution contains Fe²⁺ ions from the unrusted iron dissolving in the acid, and, Fe³⁺ ions from the rusted iron.
 - 25.0 cm³ of this solution required 16.9 cm³ of 0.020 mol dm⁻³ KMnO₄ for complete oxidation of the Fe²⁺ ions. Calculate the moles of Fe²⁺ ions in the sample titrated.
 - To a second 25.0 cm³ of the rusted iron solution an oxidising agent was added to convert all the Fe²⁺ ions present to Fe³⁺ ions. The Fe³⁺ ions were titrated with a solution of EDTA⁴⁻(aq) ions and 17.6 cm³ of 0.10 mol dm⁻³ EDTA were required. Assuming 1 mole of EDTA reacts with 1 mole of Fe³⁺ ions, calculate the moles of Fe³⁺ ions in the sample.
 - From your calculations in (a) and (b) calculate the ratio of rusted iron to unrusted iron and hence the percentage of iron that had rusted.